n. 14

Serial No. 09/981,289 Filed: October 15, 2001

Please amend the application as follows:

## In the claims:

Please cancel claims 1-11 without prejudice or disclaimer as drawn to non-elected inventions.

Please add the following new claims;

cellow cy. en 312 gcel

1/2. A

- . A method for generating a secondary library of scaffold protein variants comprising:
  - a) generating a probability distribution table of amino acid residues in a plurality of variant positions utilizing a force field calculation; and
  - b) combining a plurality of said amino acid residues to generate a secondary library of secondary sequences;

wherein at least one of said secondary variants is different from said primary variants.

13. A method according to claim 12, wherein said force field calculation is SCMF.

14. A method according to claim 12, further comprising computationally recombining said secondary library to generate a tertiary library.

15. A method according to claim 14, wherein Protein Design Automation (PDA<sup>TM</sup>) is used to recombine said secondary library.

- 16. A method for generating a secondary library of scaffold protein variants comprising:
  - a) generating a probability distribution table of amino acid residues in a plurality of variant positions utilizing an alignment program;
  - b) combining a plurality of said amino acid residues to g nerate a secondary library of secondary sequences, wherein at least one of said secondary variants is different from

Serial No. 09/981,289
- Filed: October 15, 2001

said primary variants; and

- c) computational ranking said secondary library.
- 17. A method according to claim 16, further comprising computationally recombining said secondary library to generate a tertiary library.
- 18. A method according to claim 17, wherein PDATM is used to recombine said secondary library.
- 19. A method according to claim 16, wherein said alignment program is a sequence alignment program.



- 20. A method according to claim 16, wherein said alignment program is a structural alignment program.
- 21. A method according to claims 12 or 16 further comprising synthesizing a plurality of said secondary sequences.
- 22. A method according to claim 21 wherein said synthesizing is done by multiple PCR with pooled oligonucleotides.
- 23. A method according to 22 wherein said pooled oligonucleotides are added in equimolar amounts.